
SP-TOX

A companion of Toxic Management



Mycotoxins are secondary metabolites produced by certain moulds / fungi. Mycotoxicos is causes less overt signs but often is associated with hidden losses due to immunosuppression, hepatic & renal dysfunction, poor health & productivity. Toxin reduces the organs function such as liver & kidney. It also negatively impact on the poultry immune system & native defense mechanism.

SP TOX has an enormous surface area so it adsorbs wide range of toxins in the intestine. It is a multi-component toxin binder specially developed as a complete solution to the Mycotoxins problem. Selected ingredients binds wide range of various toxins presence in the Feed.

Key features:

- SP TOX formulated with selected high quality ingredients to perform as a broad Spectrum Mycotoxin absorbent in challenge condition.
- Its largest surface area to bind extreme and various toxins in wide range.
- The unique ingredients supports to control chemical toxicity due to pesticides and heavy metals.
- Selected silicates to adsorb both polar and dipolar Mycotoxins
- Maximizes Feed conversion and Improves overall performance
- SP TOX does not absorb vitamins, Minerals and other nutrients It reduces Mortality & Secondary bacterial diseases problem
- Protects the liver against the damage and mobilizes the fat accumulated in the liver

Benefits:

- · To adsorb a wide range of mycotoxins.
- To protect and rehabilities the liver and kidney.
- To reduce immunosuppression caused by mycotoxin.
- · To Improve birds health through increased resistance to disease.
- To improve the immunity and feed intake.
- It formulated with selected high quantity ingredients to perform as a broad spectrum.
- It supports to control chemical toxicity due to pesticides and heavy metals.
- To maximize feed conversion and improves overall performance.

Composition:

Specially preserved silicates, Mannan oligosaccharide (MOS), Activated Charcoal, Organic Acids, Phytogenic derivatives and the blend of adsorbing components.

Dosage:

Layer and Broiler : 0.5 Kg to 1 Kg per ton of feed at moisture level < 15%

1 Kg to 1.5 Kgs per ton of feed at moisture level > 15%

Cattle : 0.5 Kg to 1 Kg per ton of Feed.

Not for human, Cat & Dog

Animal Feed Supplement

THE PRODUCT IS FREE FROM CHLORAMPHENICOL, NITROFURAN, ANTIBIOTICS, MELAMINE, BY-PRODUCT OF SWINE/BOUVINE AND ANY OTHER GENETICALLY MODIFIED ORGANISM.

Manufactured & marketing:



No 11, VOC Street, Ponmalaipatti, Tiruchirappalli - 620004, Tamilnadu,India. biozonaahc@gmail.com +91 9790155126 www.biozona.in